

UGT3A2 Rabbit pAb

货号: **AYP16268**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	55kDa/59kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	B cells,Jurkat,HepG2,Mouse kidney,Mouse spleen,Rat liver,Rat spleen
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 23-300 of human UGT3 A2 (NP_777574.2).
序列	AKILTISTVGGSHYLLMDRVSQILQDHGHNVTMLNHNKRGPFMPDFKKEEKSYQVISWLAPEDHQREFKKSFDFFLEETLG GRGKFENLLNVLEYLALQCSHFLNRKDIMDSLKNENFDMVIVETFDYCPFLIAEKLGKPFVAILSTSFGLSLEFGLPIPLSYVPV FRSLTDMDFWGRVKNFLMFFSFCRRQQHMQSTFDNTIKEHFTEGSRPVLSHLLKAEWFINSDFAFDFAFARPLLNTV YVGLMEKPIKVPQDLENFIAKFGDSGFVLVTL

靶点信息

研究背景	UDP-glucuronosyltransferases catalyze phase II biotransformation reactions in which lipophilic substrates are conjugated with glucuronic acid to increase water solubility and enhance excretion. They are of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds (By similarity).
------	---

基因ID	167127
基因名	UGT3A2
Swiss	Q3SY77
别名	UGT3A2

产品验证



Western blot analysis of UGT3A2 expressed in Mouse kidney,Rat liver,B cells using UGT3A2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn