

SGK3 Rabbit pAb

货号: **AYP16278**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	53kDa/57kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HL-60,THP-1,Mouse kidney
细胞定位	Cytoplasmic vesicle,Early endosome,Recycling endosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-150 of human SGK3 (NP_037389.4).
序列	MQRDHTMDYKESCPSVSISSDEHREKKRFTVYKVLVSVGRSEWVFRRYAEDKLYNTLKKQFPAMALKIPAKRIFGDN FDPDFIKQRRAGLNEFIQNLVRYPELYNHPDVRAFLQMDSPKHQSDPSEDEDERSSQKLHSTSQINILG

靶点信息

研究背景	This gene is a member of the Ser/Thr protein kinase family and encodes a phosphoprotein with a PX (pho x homology) domain. The protein phosphorylates several target proteins and has a role in neutral amino acid transport and activation of potassium and chloride channels. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.
基因ID	23678

基因名	
Swiss	Q96BR1
别名	SGK3;CISK;SGK2;SGKL

产品验证



Western blot analysis of SGK3 expressed in HL-60,THP-1,Mouse kidney using SGK3 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn