

UGT2B10 Rabbit pAb

货号: **AYP16279**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	50kDa/60kDa
实测分子量	65kDa/80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,SKOV3,U-937,Mouse liver,Mouse kidney,Rat liver
细胞定位	Endoplasmic reticulum membrane,Microsome membrane,Single-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 20-280 of human UGT2 B10 (NP_001066.1).
序列	GSCGKVLVWAAEYSLWMNMKILKELVQRGHEVTVLASSASILFDPNDSSTLKLEVYPTSLTKTEFENIIMQLVKRLSEIQK DTFWLPFSQEQEILWAINDIIRNFCKDVVSNKKLMKKLQESRFDIVFADAYLPCGELLAELFNIPFVYSHSFSPGYSEFERHSG GFIFPPSYVPVVMKLSLSDQMTFMERVKNMLYVLYFDFWFQIFNMKKWDQFYSEVLGRPTTLSETMRKADIWLMRNSWNF KFPHPFLPNVDFVGGGLH

靶点信息

研究背景	UDPGT is of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds.
基因ID	7365

基因名	UGT2B10
Swiss	P36537
别名	UGT2B10;UDPGT2B10

产品验证



Western blot analysis of UGT2B10 expressed in Mouse liver,Rat liver,293T using UGT2B10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn