

— ABLYBIO, Help Your Research



STK19 Rabbit pAb

货号: **AYP16280**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	28kDa/40kDa
实测分子量	25KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse intestine,Mouse testis,Mouse liver,Rat lung
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 75-364 of human STK19 (NP_004188.1).
------	--

靶点信息

研究背景	This gene encodes a serine/threonine kinase which localizes predominantly to the nucleus. Its specific function is unknown; it is possible that phosphorylation of this protein is involved in transcriptional regulation. This gene localizes to the major histocompatibility complex (MHC) class III region on chromosome 6 and expresses two transcript variants.
基因ID	8859
基因名	STK19
Swiss	P49842 (https://www.uniprot.org/uniprotkb/P49842/entry)
别名	STK19,D6S60,D6S60E,G11,HLA-RP1,RP1,STK19 Rabbit pAb,Inactive serine/threonine-protein kinase 19,Protein G11,Protein RP1,Tandem winged helix protein formerly known as STK19

产品验证



Western blot analysis of STK19 expressed in Mouse intestine,Rat lung,Mouse testis using STK19 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)