

RAC3 Rabbit pAb

货号: **AYP16315**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	21kDa
实测分子量	21kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BT-474,HL-60,Human placenta,Mouse lung,Mouse brain,Mouse thymus
细胞定位	Cell membrane,Cell projection,Cytoplasm,Endomembrane system,cytoskeleton,lamellipodium,perinuclear region
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-192 of human RAC3 (NP_005043.1).
序列	MQAIKCVVVG DGAVGKTCLLISYTTNAFPGEYIPTVFDNYSANVMVDGKPVNLGLWDTAGQEDYDRLRPLSYPQTDVFLI CFLVSPASFENVRAKWYPEVRHHCPTHILLVGTGLDLRDDKDTIERLRDKKLAPITYPQGLAMAREIGSVKYLECSALTQR GLKTVFDEAIRAVLCPPPVKKPGKKCTVF

靶点信息

研究背景	The protein encoded by this gene is a GTPase which belongs to the RAS superfamily of small GTP-binding proteins. Members of this superfamily appear to regulate a diverse array of cellular events, including the control of cell growth, cytoskeletal reorganization, and the activation of protein kinases. Alternative splicing results in multiple transcript variants.
------	---

基因ID	5881
基因名	RAC3
Swiss	P60763
别名	RAC3

产品验证



Western blot analysis of RAC3 expressed in Human placenta, Mouse lung, BT-474 using RAC3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn