

— ABLYBIO, Help Your Research



# RRAGC Rabbit pAb

货号: **AYP16322**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	44kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,293T,HL-60,Mouse lung,Mouse ovary
细胞定位	Cytoplasm,Lysosome,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human RRAGC (NP_071440.1).
------	---

## 靶点信息

研究背景	This gene encodes a member of the GTR/RAG GTP-binding protein family. The encoded protein is a monomeric guanine nucleotide-binding protein which forms a heterodimer with RRAGA and RRAGB and is primarily localized to the cytoplasm. The encoded protein promotes intracellular localization of the mTOR complex. Alternative splicing results in multiple transcript variants.
基因ID	64121
基因名	RRAGC
Swiss	Q9HB90 ( <a href="https://www.uniprot.org/uniprotkb/Q9HB90/entry">https://www.uniprot.org/uniprotkb/Q9HB90/entry</a> )
别名	RRAGC,GTR2,RAGC,TIB929,RRAGC Rabbit pAb,GTPase-interacting protein 2

## 产品验证



Western blot analysis of RRAGC expressed in Mouse lung, HeLa, Jurkat using RRAGC Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))