

RRAGC Rabbit pAb

货号: AYP16322

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	44kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,293T,HL-60,Mouse lung,Mouse ovary
细胞定位	Cytoplasm,Lysosome,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human RRAGC (NP_071440.1).
序列	MSLQYGAEETPLAGSYGAADSFPKDFGYGVEEEEEEEAAAAGGGVAGAGGGCGPGGADSSKPRILLMGLRRSGKSSIQ KVVVFKMSPNETLFLESTNKIYKDDISNSSFVNFQIWDFFPQMDFFDPTFDYEMIFRGTGALIYVIDAQDDYMEALTRLHIT VSKAYKVNPD MNFEVFIHKVDGLSDDHKIETQRDIHQRANDDLADAGLEKLHLSFYLTSIYDHSIFEAFSKVVQKLIPQLPT LENLLNIF

靶点信息

研究背景	This gene encodes a member of the GTR/RAG GTP-binding protein family. The encoded protein is a monomeric guanine nucleotide-binding protein which forms a heterodimer with RRAGA and RRAGB and is primarily localized to the cytoplasm. The encoded protein promotes intracellular localization of the mTOR complex. Alternative splicing results in multiple transcript variants.
------	--

基因ID	64121
基因名	RRAGC
Swiss	Q9HB90
别名	RRAGC;GTR2;RAGC;TIB929

产品验证



Western blot analysis of RRAGC expressed in Mouse lung, HeLa, Jurkat using RRAGC Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn