

— ABLYBIO, Help Your Research



MTMR14 Rabbit pAb

货号: **AYP16323**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	60kDa/66kDa/72kDa
实测分子量	72KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 335-500 of human MTM R14 (NP_001070993.1).
------	--

靶点信息

研究背景	This gene encodes a myotubularin-related protein. The encoded protein is a phosphoinositide phosphatase that specifically dephosphorylates phosphatidylinositol 3,5-biphosphate and phosphatidylinositol 3-phosphate. Mutations in this gene are correlated with autosomal dominant centronuclear myopathy. Alternative splicing results in multiple transcript variants. A pseudogene of this gene is found on chromosome 18.
基因ID	64419
基因名	MTMR14
Swiss	Q8NCE2 (https://www.uniprot.org/uniprotkb/Q8NCE2/entry)
别名	MTMR14,C3orf29,MTMR14 Rabbit pAb,HCV NS5A-transactivated protein 4 splice variant A-binding protein 1,Myotubularin-related protein 14,Phosphatidylinositol-3-phosphate phosphatase,hJumpy

产品验证



Western blot analysis of MTMR14 expressed in MCF7 using MTMR14 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)