

# PNLIP Rabbit pAb

货号: **AYP16348**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	51kDa
实测分子量	51kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 166-465 of human PNLIP (NP_000927.1).
序列	IGHSLGAHAAGEAGRRTNGTIGRITGLDPAEPCFQGTPELVRLDPSDAKFVDVIHTDGAPIVPNLGFMSQVVGHLDFFPN GGVEMPGCKKNILSQIVDIDGIWEGTRDFAACNHLRSYKYTDSIVNPDGFAGFPCASYNVFTANKCFPCPSGGCPQMGH YADRYPGKTNDVGQKFYLDTGASNFARWRYKVSVTLSGKKVTGHILVSLFGNKGNSKQYEIFKGLKPDSTHSNEFDS DVDVGDLMVKFIWYNNVINPTLPRVGASKIIVETNVGKQFNFCSPETVREEVLLTLTPC

## 靶点信息

研究背景	This gene is a member of the lipase gene family. It encodes a carboxyl esterase that hydrolyzes insoluble , emulsified triglycerides, and is essential for the efficient digestion of dietary fats. This gene is expressed specifically in the pancreas.
------	--

基因ID	5406
基因名	PNLIP
Swiss	P16233
别名	PNLIP;PL;PNLIPD;PTL

## 产品验证



Western blot analysis of PNLIP expressed in BxPC-3 using PNLIP Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)