

— ABLYBIO, Help Your Research



NAA60 Rabbit pAb

货号: **AYP16356**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	19kDa/20kDa/27kDa/28kDa
实测分子量	27KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293F
细胞定位	Golgi apparatus membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-242 of human NAA60 (NP_079121.1).
------	---

靶点信息

研究背景	This gene encodes an enzyme that localizes to the Golgi apparatus, where it transfers an acetyl group to the N-terminus of free proteins. This enzyme acts on histones, and its activity is important for chromatin assembly and chromosome integrity. Alternative splicing and the use of alternative promoters results in multiple transcript variants. The upstream promoter is located in a differentially methylated region (DMR) and undergoes imprinting; transcript variants originating from this position are expressed from the maternal allele.
基因ID	79903
基因名	NAA60
Swiss	Q9H7X0 (https://www.uniprot.org/uniprotkb/Q9H7X0/entry)
别名	NAA60,HAT4,NAT15,NatF,hNaa60,NAA60 Rabbit pAb,Histone acetyltransferase type B protein 4,N-acetyltransferase 15,N-alpha-acetyltransferase F

产品验证



Western blot analysis of NAA60 expressed in 293F using NAA60 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)