

KBTBD7 Rabbit pAb

货号: **AYP16358**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	77kDa
实测分子量	77kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,HL-60,A-549,293T,Mouse brain,Mouse kidney,Rat brain,Rat kidney
细胞定位	cytosol,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 475-684 of human KBTBD7 (NP_115514.2).
序列	YSFELIVVQNYLYAVNSKRMCLCYDPSHNMWLNCASLKRSDFQEACVFNDEIYCICDIPVMKVYNPARGEWRRISNIPLDSETHNYQIVNHDQKLLLLITSTTPQWKKNRVTVYEYDTREDQWINIGTMLGLLQFDSGFICLCARVYPSCLEPGQSFITEEDDARS ESSTEWDLDGFSSELDSESGSSSFSDDEVWVQVAPQRNAQDQGGSL

靶点信息

研究背景	The protein encoded by this gene is a transcriptional activator, having been shown to increase the transcription of activator protein-1 and serum response element. The encoded protein can also form a complex with KBTBD6 and CUL3, which regulates the ubiquitylation and degradation of TIAM1, which is a regulator of RAC1. [provided by RefSeq, Jul 2016]
基因ID	84078

基因名	KBTBD7
Swiss	Q8WVZ9
别名	KBTBD7

产品验证



Western blot analysis of KBTBD7 expressed in Mouse brain,Rat brain,MCF7 using KBTBD7 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn