

# SUPT20H Rabbit pAb

货号: **AYP16361**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	80kDa/85kDa/88kDa
实测分子量	86kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,HL-60,Mouse testis,Mouse brain,Mouse lung,Rat kidney,Rat brain
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 580-779 of human SUPT 20H (NP_001014308.2).
序列	NLSGLLPSSGGLLPNALPSAMQAASQAGVPFGLKNTSSLRPLNLLQLPGGSLIFNTLQQQQQQLSQFTPQQPQQPTTCSPO QPGEQGSSEQSTSSEQALSQAQAAVINLTGVGSFMQSQAAVLSQLGSAENRPEQSLPQQRFQLSSAFQQQQQIQQL RFLQHQMAMAAAAAQAQLHHRHTGSQSKSKMKRGTPPTPKF

## 靶点信息

研究背景	Required for MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14 activation during gastrulation. R required for down-regulation of E-cadherin during gastrulation by regulating E-cadherin protein level down stream from NCK-interacting kinase (NIK and independently of the regulation of transcription by FGF signaling and Snail (By similarity. Required for starvation-induced ATG9A trafficking during autophagy.
------	--

基因ID	55578
基因名	SUPT20H
Swiss	Q8NEM7
别名	SUPT20H;C13;C13orf19;FAM48A;FP757;P38IP;SPT20

## 产品验证



Western blot analysis of SUPT20H expressed in Mouse testis, Rat kidney, SW480 using SUPT20H Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)