

— ABLYBIO, Help Your Research



SUPT20H Rabbit pAb

货号: **AYP16361**

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IHC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 |
| 理论分子量 | 80kDa/85kDa/88kDa |
| 实测分子量 | 86kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | SW480,HL-60,Mouse testis,Mouse brain,Mouse lung,Rat kidney,Rat brain |
| 细胞定位 | Nucleus |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 580-779 of human SUPT 20H (NP_001014308.2). |
|------|---|

靶点信息

| | |
|-------|---|
| 研究背景 | Required for MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14 activation during gastrulation. Required for down-regulation of E-cadherin during gastrulation by regulating E-cadherin protein level downstream from NCK-interacting kinase (NIK and independently of the regulation of transcription by FGF signaling and Snail (By similarity. Required for starvation-induced ATG9A trafficking during autophagy. |
| 基因ID | 55578 |
| 基因名 | SUPT20H |
| Swiss | Q8NEM7 (https://www.uniprot.org/uniprotkb/Q8NEM7/entry) |
| 别名 | SUPT20H,C13,C13orf19,FAM48A,FP757,P38IP,SPT20,SUPT20H Rabbit pAb,p38-interacting protein |

产品验证



Western blot analysis of SUPT20H expressed in Mouse testis, Rat kidney, SW480 using SUPT20H Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)