

MOCS3 Rabbit pAb

货号: **AYP16365**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	49kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Jurkat,Mouse ovary,Mouse heart
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 271-460 of human MOC S3 (NP_055299.1).
序列	YSGSLLLFDALRGHFRSIRLRSRRLDCAACGERPTVTDLLDYEAFCGSSATDKCRSLQLLSPEERVSVDYKRLLD SGAFHL LLDVRPQVEVDICRLPHALHIPLKHLERRDAESLKLLKEAIWEEKQGTQEGAAVPIYVICKLGND SQAVKILQSL SAAQELD PLTVRDVVGGLMAWAAKIDGTFPQY

靶点信息

研究背景	Molybdenum cofactor (MoCo) is necessary for the function of all molybdoenzymes. The protein encoded by this gene adenylates and activates molybdopterin synthase, an enzyme required for biosynthesis of Mo Co. This gene contains no introns. A pseudogene of this gene is present on chromosome 14.
基因ID	27304

基因名	MOCS3
Swiss	O95396
别名	MOCS3;UBA4

产品验证



Western blot analysis of MOCS3 expressed in Mouse ovary, HepG2, Jurkat using MOCS3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn