

MAP4K3 Rabbit pAb

货号: **AYP16371**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	98kDa/100kDa/101kDa
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,MCF7,293T,HepG2,Mouse brain,Mouse kidney,Mouse lung,Rat heart,Rat brain
细胞定位	cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 755-894 of human MAP 4K3 (NP_003609.2).
序列	ETVNPNSTSSWFTESDTPQTNVHTVHTQLERDTILVCLDCCIKIVNLQGRKSSRKLSELTFDFQIESIVCLQDSVLAFWKH GMQGRSFRSNEVTQEISDSTRIFRLLGSDRVVLESRPTDNPTANSNLYILAGHENSY

靶点信息

研究背景	This gene encodes a member of the mitogen-activated protein kinase kinase kinase kinase family. The encoded protein activates key effectors in cell signalling, among them c-Jun. Alternatively spliced transcripts encoding multiple isoforms have been observed for this gene.
基因ID	8491

基因名	MAP4K3
Swiss	Q8IVH8
别名	MAP4K3;GLK;MAPK3;MEKK3;MEKKK3;RAB8IPL1

产品验证



Western blot analysis of MAP4K3 expressed in Mouse brain,Rat heart,SW480 using MAP4K3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn