

# NFIL3 Rabbit pAb

货号: **AYP16395**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	51kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	SW620
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 133-462 of human NFIL3 (NP_005375.2).
序列	AYAQEIQKLSNSTAVYFQDYQTSKSNVSSFVDEHEPSMVSSCISVIKHSPQSSLSDVSEVSSVEHTQESSVQGSCRSPEN KFQIIKQEPMELESYTRPRDDRGSYTASIYQNYMGNSFSGYSHSPPLLQVNRSSNSPRTSETDDGVVGKSSDGEDEQQ VPKGPIHSPVELKHVHATVVKVPEVNSSALPHKLRKAKAMQIKVEAFDNEFEATQKLSSPIDMTSKRHFLEKHSAPSMVH SSLTPFSVQVTNIQDWSLKSEHWHQKELSGKTQNSFKTGVVEMKDSGYKVS DPENLYLKQGANLSAEVVSLKRLIATQPI SASDSG

## 靶点信息

研究背景	The protein encoded by this gene is a transcriptional regulator that binds as a homodimer to activating transcription factor (ATF) sites in many cellular and viral promoters. The encoded protein represses PER1 and PER2 expression and therefore plays a role in the regulation of circadian rhythm. Three transcript variants encoding the same protein have been found for this gene.
------	--

基因ID	4783
基因名	NFIL3
Swiss	Q16649
别名	NFIL3;E4BP4;IL3BP1;NF-IL3A;NFIL3A

## 产品验证



Western blot analysis of NFIL3 expressed in SW620 using NFIL3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)