

# C10orf32 Rabbit pAb

货号: **AYP16408**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	11kDa
实测分子量	11kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	BORC complex
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-105 of human C10orf32 (NP_653192.2).
序列	MMATGTPESQARFGQSVKGLLTKVTTTCGTDVIALTKQVLKGSRSSELLGQAARNMVLQEDAILHSEDSLKMAIITTHLQYQQAQKNVEQSSDLQDLNHL

## 靶点信息

研究背景	As part of the BORC complex may play a role in lysosomes movement and localization at the cell periphery. Associated with the cytosolic face of lysosomes, the BORC complex may recruit ARL8B and couple lysosomes to microtubule plus-end-directed kinesin motor.
基因ID	119032

基因名	BORCS7
Swiss	Q96B45
别名	BORCS7;C10orf32

## 产品验证



Western blot analysis of C10orf32 expressed in human breast cancer using C10orf32 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)