

# UBE2Z Rabbit pAb

货号: **AYP16425**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	28kDa/38kDa
实测分子量	25kDa/38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BT-474,MCF7,HepG2,HeLa,Mouse pancreas,Mouse testis
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 115-354 of human UBE2Z (NP_075567.2).
序列	PPPGMFVVPDVTDMTKIHALITGPFDPYEGGFLLVFVRCPPDYPIHPPRVKLMTTGNNTVRFNPNFYRNGKVCLSILGTWT GPAWSPAQSISVLSLIQSLMTENPYHNEPGFEQERHPGDSKNYNECIRHETIRVAVCDMMEGKCPCEPLRGVMEKSFLE YYDFYEVACKDRLHLQGQTMQDPFGEKRGHFDYQSLLMRLGLIRQKVLRLHNENAEMDSDSSSSSGTETDLHGSLRV

## 靶点信息

研究背景	This gene encodes an enzyme which ubiquitinates proteins which participate in signaling pathways and a poptosis.
基因ID	65264

基因名	UBE2Z
Swiss	Q9H832
别名	UBE2Z;HOYS7;USE1

## 产品验证



Western blot analysis of UBE2Z expressed in Mouse pancreas, BT-474, MCF7 using UBE2Z Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)