

GTF2H3 Rabbit pAb

货号: **AYP16436**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:50 - 1:200
理论分子量	29kDa/34kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HL-60,HT-29,MCF-7,Jurkat
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-308 of human GTF2H3 (NP_001507.2).
序列	MVSEDELNLLVIVVDANPIIWWGKQALKESQFTLSKCIDAVMVLGNSHLFMNRSNKLAVIASHIQESRFLYPGKNGRLGD FFGDPGNPPEFNPSGSKDGKYELLTSANEVIVEEIKDLMTKSDIKGQHTETLLAGSLAKALCYIHRMNKEVKDNQEMKSRIL VIKAAEDSALQYMFNMFVIFAAQKQNILIDACVLDSDSLQACDITGGLYLKVPMPSLLQYLLWVFLPDQDQRSQLIL PPPVHVDYRAACFCHRNLEIGYVCSVCLSFICNFSPICCTCETAFKISLPPVLKAKKKLKVSA

靶点信息

研究背景	This gene encodes a member of the TFB4 family. The encoded protein is a subunit of the core-TFIID basal transcription factor and localizes to the nucleus. The encoded protein is involved in RNA transcription by RNA polymerase II and nucleotide excision repair and associates with the Cdk-activating kinase complex. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 14.
------	---

基因ID	2967
基因名	GTF2H3
Swiss	Q13889
别名	GTF2H3;BTF2;P34;TFB4;TFIIH

产品验证



Western blot analysis of GTF2H3 expressed in HL-60, HT-29, MCF-7 using GTF2H3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn