

— ABLYBIO, Help Your Research



BCL6 Rabbit pAb

货号: **AYP16445**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:100
理论分子量	72kDa/78kDa
实测分子量	86KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 350-450 of human BCL6 (NP_001124 317.1).
------	---

靶点信息

研究背景	The protein encoded by this gene is a zinc finger transcription factor and contains an N-terminal POZ domain. This protein acts as a sequence-specific repressor of transcription, and has been shown to modulate the transcription of STAT-dependent IL-4 responses of B cells. This protein can interact with a variety of POZ-containing proteins that function as transcription corepressors. This gene is found to be frequently translocated and hypermutated in diffuse large-cell lymphoma (DLCL), and may be involved in the pathogenesis of DLCL. Alternatively spliced transcript variants encoding different protein isoforms have been found for this gene.
基因ID	604
基因名	BCL6
Swiss	P41182 (https://www.uniprot.org/uniprotkb/P41182/entry)
别名	BCL6,BCL5,BCL6A,LAZ3,ZBTB27,ZNF51,BCL6 Rabbit pAb,B-cell lymphoma 5 protein,Protein LAZ-3,Zinc finger and BTB domain-containing protein 27,Zinc finger protein 51

产品验证



Western blot analysis of BCL6 expressed in Raji using BCL6 Rabbit pAb at 1:1000. Secondary antibody: HR P Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)