

# BOC Rabbit pAb

货号: **AYP16446**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	100kDa/121kDa
实测分子量	121kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,SW480,U-251MG,SKOV3,HeLa,Mouse lung,Mouse brain,Rat spinal cord
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-125 of human BOC (NP_150279.1).
序列	MLRGTMATAWRGMRPEVTLACLLLATAGCFADLNEVPQVTVQPASTVQKPGGTIVLGCVVEPPRMNVTWRLNGKELNGS DDALGVLITHGTLVITALNNHTVGRYQCVARMPAGAVASVPATVTLA

## 靶点信息

研究背景	The protein encoded by this gene is a member of the immunoglobulin/fibronectin type III repeat family. It is a component of a cell-surface receptor complex that mediates cell-cell interactions between muscle precursor cells, and promotes myogenic differentiation. Alternative splicing results in multiple transcript variants encoding different isoforms.
基因ID	91653

基因名	BOC
Swiss	Q9BWW1
别名	BOC;CDON2

## 产品验证



Western blot analysis of BOC expressed in Mouse lung,Rat spinal cord,A-431 using BOC Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)