

NCR3 Rabbit pAb

货号: **AYP16450**

产品信息

反应	Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	16kDa/17kDa/18kDa/19kDa/20kDa/21kDa
实测分子量	22KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Daudi
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 20-140 of human NCR3 (NP_667341.1).
序列	WVSQPPEIRTLEGSSAFLPCSFNASQGRILAIGSVTWFRDEVVPGKEVRNGTPEFRGRLAPLASSRFLHDHQAEIHIRDVR GHDASIYVCRVEVLGLGVGTGNGTRLVVEKEHPQLGAGTVL

靶点信息

研究背景	The protein encoded by this gene is a natural cytotoxicity receptor (NCR) that may aid NK cells in the lysis of tumor cells. The encoded protein interacts with CD3-zeta (CD247), a T-cell receptor. A single nucleotide polymorphism in the 5' untranslated region of this gene has been associated with mild malaria susceptibility. Three transcript variants encoding different isoforms have been found for this gene.
基因ID	259197

基因名	NCR3
Swiss	O14931
别名	NCR3;1C7;CD337;LY117;MALS;NKp30

产品验证



Western blot analysis of NCR3 expressed in Daudi using NCR3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn