

THADA Rabbit pAb

货号: AYP16459

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:100
理论分子量	75kDa/100kDa/105kDa/183kDa/211kDa/219kDa
实测分子量	220kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,HL-60,U-87MG,NCI-H460,Mouse lung,Mouse testis
细胞定位	
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1684-1953 of human THADA (NP_071348.3).
序列	QLVILSCEDHLPTE SRLAVVEVLTSTTPLFLTNPHPILELQDTLALWKCVLTLQSEEQAVRDAATETVTTAMSQENTCQSTE FAFCQVDASIALALAVLCDLLQQWDQLAPGLPILLGWLLGESDDL VACVESHQVEEDYLFEKAEVNFWAETLIFVKYL CKHLFCLLSKSGWRPPSPPEMLCHLQRMVSEQCHLLSQFFRELPPAAEFVKTVFETRLRIQEERTLA CLRLLAFLEGKEGEDT LVLSVWDSYAESRQLTLPRTAAC

靶点信息

研究背景	This gene is the target of 2p21 chromosomal aberrations in benign thyroid adenomas. Single nucleotide polymorphisms (SNPs) in this gene may be associated with type 2 diabetes and polycystic ovary syndrome. The encoded protein is likely involved in the death receptor pathway and apoptosis.
------	---

基因ID	63892
基因名	THADA
Swiss	Q6YHU6
别名	THADA;ARMC13;GITA

产品验证



Western blot analysis of THADA expressed in Mouse lung, MCF7, HL-60 using THADA Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn