— ABLYBIO, Help Your Research

MARCKSL1 Rabbit pAb

货号: AYP16461

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	**WB:** 1:500 - 1:2000
理论分子量	19kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BT-474,22Rv1,HT-29,NCI-H460,Mouse spleen,Rat brain
细胞定位	Cell membrane,Cytoplasm,Lipid-anchor,Membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-195 of human MARCKSL1 (NP_075385.1).

靶点信息

研究背景	This gene encodes a member of the myristoylated alanine-rich C-kinase substrate (MARCKS) family. Members of this family play a role in cytoskeletal regulation, protein kinase C signaling and calmodulin signaling. The encoded protein affects the formation of adherens junction. Alternative splicing results in multiple transcript variants. Pseudogenes of this gene are located on the long arm of chromosomes 6 and 10.
基因ID	65108
基因名	MARCKSL1
Swiss	P49006 (https://www.uniprot.org/uniprotkb/P49006/entry)
别名	MARCKSL1,F52,MACMARCKS,MLP,MLP1,MRP,MARCKSL1 Rabbit pAb,MARCKS-like protein 1,Macrophage myristoylated alanine-rich C kinase substrate

产品验证



Western blot analysis of MARCKSL1 expressed in Mouse spleen,Rat brain,BT-474 using MARCKSL1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (https://www.ablybio.cn//www.ablybio.cn)