

# MARCKSL1 Rabbit pAb

货号: **AYP16461**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	19kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BT-474,22Rv1,HT-29,NCI-H460,Mouse spleen,Rat brain
细胞定位	Cell membrane,Cytoplasm,Lipid-anchor,Membrane
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-195 of human MARCKSL1 (NP_075385.1).
序列	MGSQSSKAPRGDVTAEAAAGASPAKANGQENGHVKSNGDLSPKGEGESPPVNGTDEAAGATGDAIEPAPPSQGAEAK GEVPPKETPKKKKFSFKKPKLSGLSFKRNRKEGGDSSASSPTEEEQEQEIGACSDGTAQEGKAAATPESQEPQAK GAEASAASEEEAGPQATEPSTPSGPESGPTPASAEQNE

## 靶点信息

研究背景	This gene encodes a member of the myristoylated alanine-rich C-kinase substrate (MARCKS) family. Members of this family play a role in cytoskeletal regulation, protein kinase C signaling and calmodulin signaling. The encoded protein affects the formation of adherens junction. Alternative splicing results in multiple transcript variants. Pseudogenes of this gene are located on the long arm of chromosomes 6 and 10.
基因ID	65108

基因名	MARCKSL1
Swiss	P49006
别名	MARCKSL1;F52;MACMARCKS;MLP;MLP1;MRP

## 产品验证



Western blot analysis of MARCKSL1 expressed in Mouse spleen,Rat brain,BT-474 using MARCKSL1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 3 0ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)