

— ABLYBIO, Help Your Research



# BTNL2 Rabbit pAb

货号: **AYP16467**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:100 - 1:500
理论分子量	19kDa/27kDa/28kDa/30kDa/43kDa/50kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,BT-474,HL-60,293T,Mouse liver
细胞定位	Membrane,Single-pass type II membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 246-455 of human BTNL2 (NP_062548.1).
------	---

## 靶点信息

研究背景	This gene encodes a major histocompatibility complex, class II associated, type I transmembrane protein which belongs to the butyrophilin-like B7 family of immunoregulators. It is thought to be involved in immune surveillance, serving as a negative T-cell regulator by decreasing T-cell proliferation and cytokine release. The encoded protein contains an N-terminal signal peptide, two pairs of immunoglobulin-like domains, separated by a heptad peptide sequence, and a C-terminal transmembrane domain. Naturally occurring mutations in this gene are associated with sarcoidosis, rheumatoid arthritis, ulcerative colitis, inflammatory bowel disease, myositis, type 1 diabetes, systemic lupus erythematosus, acute coronary syndrome, and prostate cancer.
基因ID	56244
基因名	BTNL2
Swiss	Q9UIR0 ( <a href="https://www.uniprot.org/uniprotkb/Q9UIR0/entry">https://www.uniprot.org/uniprotkb/Q9UIR0/entry</a> )
别名	BTNL2,BTL-II,BTN7,HSBLMHC1,SS2,BTNL2 Rabbit pAb

## 产品验证



Western blot analysis of BTNL2 expressed in Mouse liver, HepG2, BT-474 using BTNL2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)