

NCOR1 Rabbit pAb

货号: **AYP16497**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:100
理论分子量	103kDa/258kDa/270kDa
实测分子量	270KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human NCOR1 (NP_006302.2).
序列	MSSSGYPPNQGAFSTEQSRYPHVSQYTFPNTRHQQEFVAVPDYRSSHLEVSQASQLLQQQQQLRRRPSLLSEFHGSDRPQERRTSYEPFHGSPSPVDHDSLESKRPRLEQVSDSHFQRVSAAVLPLVHPLPEGLRASADAKKDPAFGGKHEAPSSPISGQPCGDDQNASPSKLSKEELIQSMRVDREIAKVEQQIL

靶点信息

研究背景	This gene encodes a protein that mediates ligand-independent transcription repression of thyroid-hormone and retinoic-acid receptors by promoting chromatin condensation and preventing access of the transcription machinery. It is part of a complex which also includes histone deacetylases and transcriptional regulators similar to the yeast protein Sin3p. This gene is located between the Charcot-Marie-Tooth and Smith-Magenis syndrome critical regions on chromosome 17. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 17 and 20.
基因ID	9611
基因名	NCOR1
Swiss	O75376
别名	NCOR1;N-CoR;N-CoR1;PPP1R109;TRAC1;hN-CoR

产品验证



Western blot analysis of NCOR1 expressed in Jurkat using NCOR1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn