

NMT2 Rabbit pAb

货号: **AYP16500**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:100
理论分子量	56kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,293T,SW480,Mouse testis
细胞定位	Cytoplasm,Membrane,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-120 of human NMT2 (NP_004799.1).
序列	MAEDSESAASQSQSLELDDQDTCGIDGDNEEETEHAAGSPGGYLGAKKKKKQKRKKEKPNSSGGTKSDSASDSQEIKIQQ PSKNPSVPMQKLQDIQRAMELLSACQGPARNIDEAAKHRYQ

靶点信息

研究背景	This gene encodes one of two N-myristoyltransferase proteins. N-terminal myristoylation is a lipid modification that is involved in regulating the function and localization of signaling proteins. The encoded protein catalyzes the addition of a myristoyl group to the N-terminal glycine residue of many signaling proteins, including the human immunodeficiency virus type 1 (HIV-1) proteins, Gag and Nef. Alternative splicing results in multiple transcript variants.
------	--

基因ID	9397
基因名	NMT2
Swiss	O60551
别名	NMT2

产品验证



Western blot analysis of NMT2 expressed in Mouse testis, Raji, 293T using NMT2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn