

RFC2 Rabbit pAb

货号: **AYP16580**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	35kDa/39kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,22Rv1,Mouse spleen
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 38-297 of human RFC2 (NP_852136.1).
序列	WVEKYRPVKLNEIVGNEDTVSRLEVFAREGNVPNIIAGPPGTGKTTSSILCLARALLGPALKDAMLELNASNDRGIDVVRNKI KMFAQQKVTLPKGRHKIIILDEADSM TDGAQQALRRTMEIYSKTTRFALACNASDKIIEPIQSRCAVLR YTKLTDAQILTRLM NVIEKERV PYTDDGLEAIIFTAQGDMRQALNNLQSTFSGFGFINSENVFKVCDEPHLLVKEMIQHCVNANIDEAYKIL AHL WHLGYSPEDIIG

靶点信息

研究背景	This gene encodes a member of the activator 1 small subunits family. The elongation of primed DNA templates by DNA polymerase delta and epsilon requires the action of the accessory proteins, proliferating cell nuclear antigen (PCNA) and replication factor C (RFC). Replication factor C, also called activator 1, is a protein complex consisting of five distinct subunits. This gene encodes the 40 kD subunit, which has been shown to be responsible for binding ATP and may help promote cell survival. Disruption of this gene is associated with Williams syndrome. Alternatively spliced transcript variants encoding distinct isoforms have been described. A pseudogene of this gene has been defined on chromosome 2.
基因ID	5982
基因名	RFC2
Swiss	P35250
别名	RFC2;RFC40

产品验证



Western blot analysis of RFC2 expressed in LO2,22Rv1,Mouse spleen using RFC2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn