

— ABLYBIO, Help Your Research



TSC22D3 Rabbit pAb

货号: **AYP16598**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	13kDa/14kDa/22kDa
实测分子量	15kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse spleen
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-79 of human TSC22D3 (NP_001015881.1).
------	---

靶点信息

研究背景	This gene encodes the anti-inflammatory protein glucocorticoid (GC)-induced leucine zipper. Expression of this gene stimulated by glucocorticoids and interleukin 10 and it appears to play a key role in the anti-inflammatory and immunosuppressive effects of this steroid. This protein has also been shown to inhibit pro-inflammatory molecules including nuclear factor κ B. Alternate splicing results in multiple transcript variants.
基因ID	1831
基因名	TSC22D3
Swiss	Q99576 (https://www.uniprot.org/uniprotkb/Q99576/entry)
别名	TSC22D3,DIP,DSIPI,GILZ,TSC-22R,TSC22D3 Rabbit pAb,DSIP-immunoreactive peptide,Delta sleep-inducing peptide immunoreactor,Glucocorticoid-induced leucine zipper protein,TSC-22-like protein,TSC-22-related protein

产品验证



Western blot analysis of TSC22D3 expressed in 293T, Mouse spleen using TSC22D3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)