

— ABLYBIO, Help Your Research



TDRKH Rabbit pAb

货号: **AYP16605**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	57kDa/62kDa
实测分子量	72kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HepG2,MCF7,SKOV3,Mouse testis,Rat testis
细胞定位	Cytoplasm,Mitochondrion
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 36-355 of human TDRKH (NP_001077432.1).
------	---

靶点信息

研究背景	Participates in the primary piRNA biogenesis pathway and is required during spermatogenesis to repress transposable elements and prevent their mobilization, which is essential for the germline integrity. The piRNA metabolic process mediates the repression of transposable elements during meiosis by forming complexes composed of piRNAs and Piwi proteins and govern the methylation and subsequent repression of transposons. Required for the final steps of primary piRNA biogenesis by participating in the processing of 31-37 nt intermediates into mature piRNAs. May act in pi-bodies and piP-bodies by transferring piRNA precursors or intermediates to or between these granules.
基因ID	11022
基因名	TDRKH
Swiss	Q9Y2W6 (https://www.uniprot.org/uniprotkb/Q9Y2W6/entry)
别名	TDRKH,TDRD2,TDRKH Rabbit pAb,Tudor domain-containing protein 2

产品验证



Western blot analysis of TDRKH expressed in Mouse testis,Rat testis,293T using TDRKH Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)