

STRN3 Rabbit pAb

货号: **AYP16608**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:100 - 1:200 IF/ICC: 1:50 - 1:100
理论分子量	77kDa/87kDa
实测分子量	87kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,Mouse kidney
细胞定位	Cytoplasm,Membrane,Peripheral membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 200-460 of human STRN 3 (NP_001077362.1).
序列	VRSLLGLSNSEPNQSVETKNLEQILNGGESPKQKQGEIKRSSGDVLETFNFLENADDSDEDEENDMIEGIPEGKDKHRMN KHKIGNEGLAADLTDDPDTEELKEFDLVTAEDEGEGAGEARSSGDGTEWDKDDLSPTAEVWDVDQGLISKLEQYKKE RKGKKGVKRANRTKLYDMIADLGDELPHIPSGIINQSRSASTRMTDHEGARAEAEPIITFPSSGGGKSFIMGSDDVLLSVL GLGDLADLTVTNDADYSYDLP

靶点信息

研究背景	Binds calmodulin in a calcium dependent manner. May function as scaffolding or signaling protein.
基因ID	29966

基因名	STRN3
Swiss	Q13033
别名	STRN3;PPP2R6B;S/G2NA;SG2NA

产品验证



Western blot analysis of STRN3 expressed in SH-SY5Y, Mouse kidney using STRN3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn