

# SPINT2 Rabbit pAb

货号: **AYP16609**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:100
理论分子量	21kDa/28kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 28-197 of human SPINT 2 (NP_066925.1).
序列	ADRERSIHDFCLVSKVVGRCRASMPRWYVNVTDGSCQLFVYGGCDGNSNNYLTKEECLKKCATVTENATGDLATSRNA ADSSVPSAPRRQDSEDHSSDMFNYYEYCTANAVTGPCRASFPFRWYFDVERNSCNNFIYGGCRGNKNSYRSEEACMLRC FRQQENPPLPLGSK

## 靶点信息

研究背景	This gene encodes a transmembrane protein with two extracellular Kunitz domains that inhibits a variety of serine proteases. The protein inhibits HGF activator which prevents the formation of active hepatocyte growth factor. This gene is a putative tumor suppressor, and mutations in this gene result in congenital sodium diarrhea. Multiple transcript variants encoding different isoforms have been found for this gene.
------	---

基因ID	10653
基因名	SPINT2
Swiss	O43291
别名	SPINT2;DIAR3;HAI-2;HAI2;Kop;PB

## 产品验证



Western blot analysis of SPINT2 expressed in MCF7 using SPINT2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)