

RBP2 Rabbit pAb

货号: **AYP16620**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:100
理论分子量	15kDa
实测分子量	15-17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse small intestine,Rat liver
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-134 of human RBP2 (NP_004155.2).
序列	MTRDQNGTWEMESNENFEGYMKALDIDFATRKIIVRLTQTKVIDQDGNFKTKTTSTFRNYDVFDFTVGVFDEYTKSLD NRHVKALVTWEGDVLVLCVQKGEKENRGWKQWIEGDKLYLELTCDGQVCRQVFKKK

靶点信息

研究背景	This gene encodes an abundant protein present in the small intestinal epithelium. It is thought to participate in the uptake and/or intracellular metabolism of vitamin A. Vitamin A is a fat-soluble vitamin necessary for growth, reproduction, differentiation of epithelial tissues, and vision. This protein may also modulate the supply of retinoic acid to the nuclei of endometrial cells during the menstrual cycle.
基因ID	5948

基因名	RBP2
Swiss	P50120
别名	RBP2;CRABP-II;CRBP2;CRBP2II;RBPC2

产品验证



Western blot analysis of RBP2 expressed in Mouse small intestine,Rat liver using RBP2 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn