

— ABLYBIO, Help Your Research



# RECK Rabbit pAb

货号: **AYP16621**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	106kDa
实测分子量	106kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,MCF-7,H460,Mouse lung,Mouse kidney
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 110-210 of human RECK (NP_066934.1).
------	--

## 靶点信息

研究背景	The protein encoded by this gene is a cysteine-rich, extracellular protein with protease inhibitor-like domains whose expression is suppressed strongly in many tumors and cells transformed by various kinds of oncogenes. In normal cells, this membrane-anchored glycoprotein may serve as a negative regulator for matrix metalloproteinase-9, a key enzyme involved in tumor invasion and metastasis. Several transcript variants encoding different isoforms have been found for this gene.
基因ID	8434
基因名	RECK
Swiss	O95980 ( <a href="https://www.uniprot.org/uniprotkb/O95980/entry">https://www.uniprot.org/uniprotkb/O95980/entry</a> )
别名	RECK,ST15,RECK Rabbit pAb,Suppressor of tumorigenicity 15 protein

## 产品验证



Western blot analysis of RECK expressed in Mouse lung,293T,MCF-7 using RECK Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))