

— ABLYBIO, Help Your Research



NPHP1 Rabbit pAb

货号: **AYP16634**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	69kDa/77kDa/83kDa
实测分子量	83KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	PC-3,Mouse pancreas,Rat skeletal muscle,Rat brain
细胞定位	Cell junction,Cell projection,Cytoplasm,adherens junction,cilium,cilium axoneme,cytoskeleton,tight junction
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 345-614 of human NPHP 1 (NP_001121651.1).
------	---

靶点信息

研究背景	This gene encodes a protein with src homology domain 3 (SH3) patterns. This protein interacts with Crk-associated substrate, and it appears to function in the control of cell division, as well as in cell-cell and cell-matrix adhesion signaling, likely as part of a multifunctional complex localized in actin- and microtubule-based structures. Mutations in this gene cause familial juvenile nephronophthisis type 1, a kidney disorder involving both tubules and glomeruli. Defects in this gene are also associated with Senior-Loken syndrome type 1, also referred to as juvenile nephronophthisis with Leber amaurosis, which is characterized by kidney and eye disease, and with Joubert syndrome type 4, which is characterized by cerebellar ataxia, oculomotor apraxia, psychomotor delay and neonatal breathing abnormalities, sometimes including retinal dystrophy and renal disease. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	4867
基因名	NPHP1
Swiss	O15259 (https://www.uniprot.org/uniprotkb/O15259/entry)
别名	NPHP1,JBTS4,NPH1,SLSN1,NPHP1 Rabbit pAb,Juvenile nephronophthisis 1 protein

产品验证



Western blot analysis of NPHP1 expressed in Mouse pancreas, Rat skeletal muscle, PC-3 using NPHP1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)