

# GLUD2 Rabbit pAb

货号: **AYP16664**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	<a href="#">WB</a> <a href="#">IHC</a> <a href="#">IF/ICC</a>
推荐浓度	<b>WB:</b> 1:100 - 1:500 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	61kDa
实测分子量	52KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,A-431,U-937
细胞定位	Mitochondrion matrix
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 319-558 of human GLU D2 (NP_036216.2).
序列	YLHRFGAKCIAVGESDGSIWNPDGIDPKELEDFKLQHGSILGFPKAKPYEGSILEVDCDILIPAATEKQLTKSNAPRVKAKIIA EGANGPTTPEADKIFLERNILVIPDLYLNAGGVTVSYFEWLKLNHVSYGRLTFKYERDSNYHLLLSVQESLERKFGKHGGTI PIVPTAEFQDSISGASEKDIVHSALAYTMERSARQIMHTAMKYNLGLDLRRTAAYVNAIEKVFKVYSEAGVTFT

## 靶点信息

研究背景	The protein encoded by this gene is localized to the mitochondrion and acts as a homohexamer to recycle glutamate during neurotransmission. The encoded enzyme catalyzes the reversible oxidative deamination of glutamate to alpha-ketoglutarate. This gene is intronless.
------	---

基因ID	2747
基因名	GLUD2
Swiss	P49448
别名	GLUD2;GDH2;GLUDP1

## 产品验证



Western blot analysis of GLUD2 expressed in HepG2,A-431,U-937 using GLUD2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)