

— ABLYBIO, Help Your Research



EIF2S3 Rabbit pAb

货号: **AYP16671**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200
理论分子量	51kDa
实测分子量	51KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HepG2,Raji,U-251MG,Mouse liver,Mouse thymus,Rat liver
细胞定位	cytoplasm,cytosol,eukaryotic translation initiation factor 2 complex,extracellular exosome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human EIF2S3 (NP_001406.1).
------	--

靶点信息

研究背景	The protein encoded by this gene is the largest subunit of a heterotrimeric GTP-binding protein involved in the recruitment of methionyl-tRNA(i) to the 40 S ribosomal subunit.
基因ID	1968
基因名	EIF2S3
Swiss	P41091 (https://www.uniprot.org/uniprotkb/P41091/entry)
别名	EIF2S3,EIF2,EIF2G,EIF2gamma,MRXSBRK,eIF-2gA,EIF2S3 Rabbit pAb,Eukaryotic translation initiation factor 2 subunit gamma X

产品验证



Western blot analysis of EIF2S3 expressed in Mouse liver,Rat liver,HeLa using EIF2S3 Rabbit pAb at 1:100. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)