

CD200 Rabbit pAb

货号: **AYP16682**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	30kDa/31kDa/32kDa
实测分子量	45-50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Rat brain
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-240 of human CD200 (NP_005935.4).
序列	AQVQVVTQDEREQLYTPASLKCSLQNAQEALIVTWQKKKAVSPENMVTFSENHGVVIQPAYKDKINITQLGLQNSTITFWN ITLEDEGCYMCLFNTFGFGKISGTACLTIVYVQPIVSLHYKFSEDHLNITCSATARPAPMVFWKVPRSGIENSTVTLSHPNGTT SVTSILHIKDPKNQVGKEVICQLHLGTVTDFKQTVNKGWFSVPLL

靶点信息

研究背景	This gene encodes a type I membrane glycoprotein containing two extracellular immunoglobulin domains , a transmembrane and a cytoplasmic domain. This gene is expressed by various cell types, including B cells, a subset of T cells, thymocytes, endothelial cells, and neurons. The encoded protein plays an important role in immunosuppression and regulation of anti-tumor activity. Alternative splicing results in multiple transcript variants encoding different isoforms.
------	--

基因ID	4345
基因名	CD200
Swiss	P41217
别名	CD200;MOX1;MOX2;MRC;OX-2

产品验证



Western blot analysis of CD200 expressed in Mouse brain,Rat brain using CD200 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn