

# AMHR2 Rabbit pAb

货号: **AYP16702**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	51kDa/52kDa/62kDa
实测分子量	58-63kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung,Rat kidney
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 18-149 of human AMHR 2 (NP_065434.1).
序列	PPNRRTCvFFEAPGVRGStKTLGELLDTGTelpRAIRCLYSRCCFGIWNLTQDRAQVEMQGRDSDEPGCESLHCDPSRAHPSPGStLFTcscGtDFCNANYSHLPPPGSPGTPGSQGPQAAPGESIWMAL

## 靶点信息

研究背景	This gene encodes the receptor for the anti-Mullerian hormone (AMH) which, in addition to testosterone, results in male sex differentiation. AMH and testosterone are produced in the testes by different cells and have different effects. Testosterone promotes the development of male genitalia while the binding of AMH to the encoded receptor prevents the development of the mullerian ducts into uterus and Fallopian tubes. Mutations in this gene are associated with persistent Mullerian duct syndrome type II. Alternatively spliced transcript variants encoding different isoforms have been identified.
------	--

基因ID	269
基因名	AMHR2
Swiss	Q16671
别名	AMHR2;AMHR;MISR2;MISRII;MRII

## 产品验证



Western blot analysis of AMHR2 expressed in Mouse lung,Rat kidney using AMHR2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)