

ACTR1A Rabbit pAb

货号: **AYP16705**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	42kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,Mouse skeletal muscle,Mouse spleen,Mouse lung,Rat brain
细胞定位	Cytoplasm,centrosome,cytoskeleton,microtubule organizing center
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 187-376 of human ACTR1A (NP_005727.1).
序列	GRDVSRLFRLYLRLKEGYDFHSSEFEIVKAIKERACYLSINPQKDETLETEKAQYYLPDGSSTIEIGPSRFRAPPELLFRPDLIGEE SEGIHEVLVFAIQSDMDLRRTLFSNIVLSGGSTLFKGFGRLLSEVKKLAPKDVKIRISAPQERLYSTWIGGSILASLDTFKK MWVSKKEYEEDGARSIHRTKTF

靶点信息

研究背景	This gene encodes a 42.6 kD subunit of dynactin, a macromolecular complex consisting of 10-11 subunits ranging in size from 22 to 150 kD. Dynactin binds to both microtubules and cytoplasmic dynein. It is involved in a diverse array of cellular functions, including ER-to-Golgi transport, the centripetal movement of lysosomes and endosomes, spindle formation, chromosome movement, nuclear positioning, and axonogenesis. This subunit is present in 8-13 copies per dynactin molecule, and is the most abundant molecule in the dynactin complex. It is an actin-related protein, and is approximately 60% identical at the amino acid level to conventional actin.
基因ID	10121
基因名	ACTR1A
Swiss	P61163
别名	ACTR1A;ARP1;Arp1A;CTRN1

产品验证



Western blot analysis of ACTR1A expressed in Mouse skeletal muscle, Rat brain, MCF7 using ACTR1A Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn