

FLCN Rabbit pAb

货号: **AYP16707**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:100
理论分子量	20kDa/37kDa/64kDa
实测分子量	70KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,A-431
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 341-579 of human FLCN (NP_659434.2).
序列	RKLPVFKSLRHMRQVLGAPSFRLAWHVLGMGNQVIWKS RDVDLVQSAFEVLR TMLPVGCVRIIPYSSQYEEAYRCN FLGLSPHVQIPPHVLSSEFAVIVEVHAAARSTLHPVGCEDDQSLSKYEFVVTSGSPVAADR VGPTILNKIEAALTNQNLSVDVVDQCLVCLKEEWMNKVKVLFKFTKVDSRPKEDTQKLLSILGASEEDNVKLLKFWMTGLSKTYKSHLMSTVRSPTASESRN

靶点信息

研究背景	This gene is located within the Smith-Magenis syndrome region on chromosome 17. Mutations in this gene are associated with Birt-Hogg-Dube syndrome, which is characterized by fibrofolliculomas, renal tumors, lung cysts, and pneumothorax. Alternative splicing of this gene results in two transcript variants encoding different isoforms.
------	--

基因ID	201163
基因名	FLCN
Swiss	Q8NFG4
别名	FLCN;BHD;FLCL

产品验证



Western blot analysis of FLCN expressed in HeLa,A-431 using FLCN Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn