

TPST2 Rabbit pAb

货号: **AYP16714**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:100 - 1:500
理论分子量	41kDa
实测分子量	42KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U2OS,Mouse spleen,Rat spleen
细胞定位	Golgi apparatus membrane,Single-pass type II membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 280-370 of human TPST 2 (NP_003586.3).
序列	SKIERSTDQVIKPVNLEALSKWTGHIPGDVVRDMAQIAPMLAQLGYDPYANPPNYGNPDPFVINNTQRVLKGDYKTPANLK GYFQVNQNST

靶点信息

研究背景	The protein encoded by this gene catalyzes the O-sulfation of tyrosine residues within acidic regions of proteins. The encoded protein is a type II integral membrane protein found in the Golgi body. Two transcript variants encoding the same protein have been found for this gene.
基因ID	8459

基因名	TPST2
Swiss	O60704
别名	TPST2;TANGO13B

产品验证



Western blot analysis of TPST2 expressed in U2OS, Mouse spleen, Rat spleen using TPST2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn