

NRBF2 Rabbit pAb

货号: AYP16720

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC IHC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200 IHC: 1:50 - 1:200
理论分子量	26kDa/31kDa/32kDa
实测分子量	38KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	Cytoplasm,Cytoplasmic vesicle,Nucleus,autophagosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-287 of human NRBF2 (NP_110386.2).
序列	MEVMEGPLNLAHQSSRRADRLLAAGKYEEAISCHKKAAAYLSEAMKLTQSEQAHLSELELQRDASHMKQLLLIQRWKRAQR EERLKAQNTDKDAAAHLQTSHKPSAEDAEGQSPLSQKYPSTEKCLPEIQGIFDRDPDTLLYLLQQKSEPAEPCIGSKAP KDDKTIIEEQATKIADLKRHVEFLVAENERLRKENKQLKAEKARLLKGPIEKELDVDADFVETSELWSLPPHAETATASSTW QKFAANTGKAKDIPIPNLPLDFPSPPELPLMELSEDILKGFMMN

靶点信息

研究背景	May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (i n vitro).
------	--

基因ID	29982
基因名	NRBF2
Swiss	Q96F24
别名	NRBF2;COPR;COPR1;COPR2;NRBF-2

产品验证



Western blot analysis of NRBF2 expressed in HepG2 using NRBF2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn