

— ABLYBIO, Help Your Research



# NRBF2 Rabbit pAb

货号: **AYP16720**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC IHC
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IF/ICC:</b> 1:50 - 1:200 <b>IHC:</b> 1:50 - 1:200
理论分子量	26kDa/31kDa/32kDa
实测分子量	38KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	Cytoplasm,Cytoplasmic vesicle,Nucleus,autophagosome
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-287 of human NRBF2 (NP_110386.2).
------	-------------------------------------------------------------------------------------------------------------------

## 靶点信息

研究背景	May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).
基因ID	29982
基因名	NRBF2
Swiss	Q96F24 ( <a href="https://www.uniprot.org/uniprotkb/Q96F24/entry">https://www.uniprot.org/uniprotkb/Q96F24/entry</a> )
别名	NRBF2,COPR,COPR1,COPR2,NRBF-2,NRBF2 Rabbit pAb,Comodulator of PPAR and RXR

## 产品验证



Western blot analysis of NRBF2 expressed in HepG2 using NRBF2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)