

— ABLYBIO, Help Your Research



CLIC1 Rabbit pAb

货号: **AYP16770**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	26kDa
实测分子量	27kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,Jurkat,HeLa,HepG2,Mouse kidney,Mouse lung,Rat kidney
细胞定位	Cell membrane,Cytoplasm,Nucleus,Nucleus membrane,Single-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 42-241 of human CLIC1 (NP_001279.2).
------	--

靶点信息

研究背景	Chloride channels are a diverse group of proteins that regulate fundamental cellular processes including stabilization of cell membrane potential, transepithelial transport, maintenance of intracellular pH, and regulation of cell volume. Chloride intracellular channel 1 is a member of the p64 family; the protein localizes principally to the cell nucleus and exhibits both nuclear and plasma membrane chloride ion channel activity.
基因ID	1192
基因名	CLIC1
Swiss	O00299 (https://www.uniprot.org/uniprotkb/O00299/entry)
别名	CLIC1,CL1C1,G6,NCC27,CLIC1 Rabbit pAb,Chloride channel ABP,Glutaredoxin-like oxidoreductase CLIC1,Glutathione-dependent dehydroascorbate reductase CLIC1,Nuclear chloride ion channel 27,Regulatory nuclear chloride ion channel protein

产品验证



Western blot analysis of CLIC1 expressed in Mouse kidney,Rat kidney,MCF7 using CLIC1 Rabbit pAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)