

RHOH Rabbit pAb

货号: **AYP16772**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:10 - 1:100
理论分子量	21kDa
实测分子量	27kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BT-474,SW620,Jurkat,HT-29,Mouse spleen,Mouse thymus,Mouse liver
细胞定位	Cell membrane,Cytoplasm,Cytoplasmic side,Lipid-anchor
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-191 of human RHOH (NP_004301.1).
序列	MLSSIKCVLVGDSAVGKTSLLVRFVTSFPEAYKPTVYENTGVDVFMDSIQISLGLWDTAGNDAFRSIRPLSYQQADVLM CYSVANHNSFLNLKNKWIGEIRSNLPCTPVLVVATQTDQREMGPHRASCVNAMEGKLAQDVRAGYLECSALSNRGVQ QVFECVAVRTAVNQARRRRRRRLFSINECKIF

靶点信息

研究背景	The protein encoded by this gene is a member of the Ras superfamily of guanosine triphosphate (GTP)-metabolizing enzymes. The encoded protein is expressed in hematopoietic cells, where it functions as a negative regulator of cell growth and survival. This gene may be hypermutated or misexpressed in leukemias and lymphomas. Chromosomal translocations in non-Hodgkin's lymphoma occur between this locus and B-cell CLL/lymphoma 6 (BCL6) on chromosome 3, leading to the production of fusion transcripts. Alternative splicing in the 5' untranslated region results in multiple transcript variants that encode the same protein.
基因ID	399
基因名	RHOH
Swiss	Q15669
别名	RHOH;ARHH;TTF

产品验证



Western blot analysis of RHOH expressed in Mouse spleen, BT-474, SW620 using RHOH Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn