

# JADE1 Rabbit pAb

货号: AYP16776

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:10 - 1:100
理论分子量	58kDa/94kDa/95kDa
实测分子量	108kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Mouse liver
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 210-509 of human JADE 1 (NP_079176.2).
序列	QSPDGEDGNEMVFCDKCNICVHQACYGILKVPEGSWLCRTCALGVQPKLLCPKGGAMKPTRSGTKWVHVSCALWIPE VSIGSPEKMEPITKVSHIPSSRWALVCSLCNEKFGASIQCSVKNCRTAFHVTCADFDRGLEMKTILAENDEVKFKSYCPKHSS HRKPEESLGKGAQENGAPECSRNPLEPFASLEQNREEAHRVSVRKQKLQLEDEFYTFVNLDDVARALRLPEEVDFLY QYWKLKRKVNFNKPLITPKKDEEDNLAKREQDVLFRRLQLFTHLRQDLERVMIDTDTL

## 靶点信息

研究背景	Scaffold subunit of some HBO1 complexes, which have a histone H4 acetyltransferase activity. Plays a key role in HBO1 complex by directing KAT7/HBO1 specificity towards histone H4 acetylation (H4K5ac, H4K8ac and H4K12ac, regulating DNA replication initiation, regulating DNA replication initiation. May also promote acetylation of nucleosomal histone H4 by KAT5. Promotes apoptosis. May act as a renal tumor suppressor. Negatively regulates canonical Wnt signaling; at least in part, cooperates with NPHP4 in this function.
基因ID	79960
基因名	JADE1
Swiss	Q6IE81
别名	JADE1;PHF17

## 产品验证



Western blot analysis of JADE1 expressed in Mouse kidney, Mouse liver using JADE1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)