

— ABLYBIO, Help Your Research



# AGFG1 Rabbit pAb

货号: **AYP16797**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	54kDa/58kDa/60kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,HepG2,Mouse lung,Mouse testis
细胞定位	Cytoplasmic vesicle,Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human AGFG1 (NP_004495.2).
------	---

## 靶点信息

研究背景	The protein encoded by this gene is related to nucleoporins, a class of proteins that mediate nucleocytoplasmic transport. The encoded protein binds the activation domain of the human immunodeficiency virus Rev protein when Rev is assembled onto its RNA target, and is required for the nuclear export of Rev-directed RNAs. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	3267
基因名	AGFG1
Swiss	P52594 ( <a href="https://www.uniprot.org/uniprotkb/P52594/entry">https://www.uniprot.org/uniprotkb/P52594/entry</a> )
别名	AGFG1,HRB,RAB,RIP,AGFG1 Rabbit pAb,HIV-1 Rev-binding protein,Nucleoporin-like protein RIP,Rev-interacting protein,Rev/Rex activation domain-binding protein

## 产品验证



Western blot analysis of AGFG1 expressed in Mouse lung, Jurkat, HepG2 using AGFG1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)