

# HSD17B13 Rabbit pAb

货号: **AYP16803**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:1000 <b>IHC:</b> 1:50 - 1:200 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	29kDa/33kDa
实测分子量	34KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Mouse lung
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 30-211 of human HSD17B13 (NP_835236.2).
序列	RRKSVAGEIVLITGAGHGIGRQTTYEFAKRQSI LVLWDINKRGVEETA AECRKLGVTAHAYVVDCSNREEIYRSLNQVKKEV GDVTIVVNNAGTVYPADLLSTKDEEITKTFEVN I LGHFWITKALLPMMERNHGHIVTVASVCGHEGIPYLPYC SSKFAAVG FHRGLTSELQALGKTGI

## 靶点信息

研究背景	Predicted to enable oxidoreductase activity, acting on the CH-OH group of donors, NAD or NADP as acceptor and steroid dehydrogenase activity. Acts upstream of or within positive regulation of lipid biosynthetic process. Located in lipid droplet.
------	---

基因ID	345275
基因名	HSD17B13
Swiss	Q7Z5P4
别名	HSD17B13;HMFN0376;NIIL497;SCDR9;SDR16C3

## 产品验证



Western blot analysis of HSD17B13 expressed in HepG2, Mouse lung using HSD17B13 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)